

Original Article

Comparison of Mental Health Status in Mothers of Primary School Children with Attention Deficit / Hyperactivity Disorder and Mothers of Normal Children in Yazd City (2015-2016)

Ali Jafari Nodoushan¹, Fariborz Dortaj², Reza Jafari Nodoushan^{3*}

- 1- Department of educational psychology, International Imam Reza University, Mashhad, Iran.
- 2- Faculty of Psychology & Education, Allameh Tabataba'i University, Tehran, Iran.
- 3- Department of Occupational Hygiene, Faculty of Public Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Received: 2016/09/05

Accepted: 2016/11/24

Abstract

Introduction: ADHD is one of the most common disorders among school children throughout the world. Parents of these children are faced with more conflicts than normal children's parents. The Purpose of his study was to evaluate and compare the mental health status of mothers having children with attention deficit / hyperactivity disorder versus mental health of mothers having normal primary school children, Yazd, Iran.

Materials and Methods: The sample consisted of 160 mothers of primary-school children who were selected through random cluster sampling; 80 of them had children with attention deficit / hyperactivity disorder and the remaining half had normal children. Also, for the diagnosis of children with attention deficit / hyperactivity disorder, the Conners test as well as test of General Mental Health (GHQ) were used to measure mothers' mental health. The data were then analyzed in two levels of descriptive and inferential statistics (T-test and analysis of variance)

Results: Comparison of mental health and its subscales indicated that mothers of children with ADHD disorder were lower in all aspects of mental health than mothers of normal children.

Conclusion: According to the research results, mothers of children with attention deficit/hyperactive disorder have lower levels of mental health than mothers of normal children. So, it is recommended that education and health officials provide training courses for these parents to promote their mental health status and consequently their quality of family life.

Keywords: Mental Health, Mothers, ADHD children

* **Corresponding author:** Tel: :+3531492204 **email:** jafarinodoushan@gmail.com

Introduction

Attention deficit / hyperactivity disorder is one of the most common neurodevelopmental disorders^[1-3] that is influenced by the genetic and environmental factors.^[4,5]

At the moment, this disorder has an increasing trend worldwide, i.e., in the United States of America 11% of children were diagnosed to have this disorder^[6]. Symptoms of this disorder are attention deficit, impulsivity, and hyperactivity^[7-9] which normally continue by adolescence^[10]. Attention deficit / hyperactivity disorder is due to the defects in executive functions^[11, 12] which includes concepts of self- monitoring, self-control, and self-regulation in children^[13, 14]. These disorders will affect children's academic performance as well as, social behavior.^[15-18]

This problem may cause a basis for children's isolation at home, school, or in relationship with peers.^[19, 20] Existence of this disorder affects functionality of the family tremendously, especially mothers' mental health^[21], it will also cast a negative impact on the relationship between the child and other members of family^[22], moreover, this situation will increase conflicts in the parents' relationship to a large extent^[23]. Different researches have presented that parents with this type of children have more stress and conflict in their relationships^[24]. Since in our country mothers spend more time with children, they are more exposed to mental damages, thus their mental health is in a threatening situation^[25]. Usually the first

treatment prescribed for such disorders is medication^[26], while drug treatments by itself will not be the only answer for this problem^[27]. Some researchers stated that parents' training about this type of children may lead to reduce the symptoms of ADHD disorder^[28, 29] and improve their methods of parenting style^[30]. In some other studies, it was indicated that parents and mothers of children with attention deficit / hyperactivity disorder are more prone to psychiatric stress and damage^[31-34]. According to the span of disorder. Role of mothers' mental health in training children and their position in Iranian families as the main foundation which keeps the balance of family structure is prominent^[35]. Therefore, evaluating and investigating mental health status of mothers with attention deficit / hyperactivity disorder children versus mothers of normal children is crucial.

Materials and Methods

This causal-comparative study consisted of 160 people selected by random cluster sampling. Half of the research populations (i.e., 80 participants) were mothers of children with ADHD disorder and the other half were mothers with normal primary school children. These groups were selected from elementary schools of the first educational regain. Initially, 6 elementary boys' schools and 3 elementary girls' schools were selected, from each school one class was chosen randomly, and then teachers of these classes were asked about any evidences of attention deficit /

hyperactivity disorder among the students. Parents also were required to answer the Conner's form to identify whether their children have any disorders. This questionnaire consisting of 27 questions was applied to diagnose attention deficit / hyperactivity disorder. The participants were asked to rate each item with a score from 0 to 3, the total scores of each questionnaire was then added and divided by the number of items. Moreover, mothers filled out and completed the general mental health questionnaires. The cut-off point in this questionnaire for this type of disorder is 1.5 and after confirmation of this condition for a child, a clinical interview was conducted with his/her parents to diagnose the disorder. The age range of 3 to 17 was set using Conners measurement. Internal correlation coefficients of durability with amplitude of 75% to 90% and Test-retest correlation coefficient with 8 distances of 60% to 90% has been reported [36].

This questionnaire has justifiability content validity and the translation copy of Tehranidoost and Zargarinezhad has been used

[36, 37]. Also, the General Health Questionnaire (GHQ-28) with the following four subscales was administered in this study:

- 1- Somatic symptoms (questions from 1 to 7)
- 2- Anxiety and insomnia (questions from 8 to 14)
- 3- Impairment of social functioning (questions 15 to 21)
- 4- Depression (questions 22 to 28)

The average time to execute this questionnaire is estimated to be between 10 to 12 minutes. For each individual, 5 scores were obtained, 4 of which were related to the subscales and 1 number came from adding up of partial measurements that finally will be the total number (i.e., 38).

Scoring method: Choosing options of A to D belongs to number from zero to three. In conclusion, by applying Table 1, each individual's situation in any sub subscales and generally in the entire questionnaire can be determined.

Table 1- Specification of Public Health Questionnaire's items in each subscale

Partial subscales	Numbers in subscales	Item Numbers
Nothing or with minimum limit	6-0	22-0
Light	11-7	40-23
Average	16-12	60-41
Intense	21-17	84-61

Reliability coefficients for this test and the subscales are respectively 0.88 and from 0.50 to 0.81, the concurrent validity coefficient is also 0.69^[37].

Result

For this research, 160 individuals were selected, 80 of whom (50%) were mothers of normal children and the other 80 (50%) were

mothers of children with attention deficit /hyperactivity disorders. Findings on grade scales and knowledge of learning are shown in tables 2 and 3, respectively. Also, descriptive findings related to age and mental health tests are presented in table 4 and results of T-TEST conducted to compare two independent groups are tabulated in Table 5.

Table 2- Frequency and percentage of educational levels of children

Base of education	frequency	percentage	real percentage	Cumulative percentage
First grade	55	34.4	34.4	34.4
Second grade	41	25.6	25.6	60
Third grade	30	18.8	18.8	78.8
Fourth grade	17	10.6	10.6	89.4
Fifth grade	9	5.6	5.6	95
Sixth grade	8	5	5	100
Total	160	100	100	

Based on Table 2, majority of children under investigation were at the first grade following by the second and third. In other words, by the higher levels of education the frequency of children lowered, until in the fifth and sixth

grades the frequency reached 21%. Results of Chi-square test that presented frequency of studied children in different bases of education were different and diverse ($\chi^2 = 66.5$).

Table 3- Frequency and percentage of educational levels of participants

Test subjects education	frequency	percentage	Cumulative percentage
Up to ninth grade(cycle)	40	25	25
Diploma	60	37.5	62.5
Over diploma	60	37.5	100
Total	160	100	

According to Table 3, the highest percentage of participants (75%) had diplomas or higher educational levels. Mother with the educational levels up to the ninth grade (25%) took the next place

Table 4- presents mothers' average and deviation of age-scales, psychology health levels and subscales of them are also represented for the two groups of mothers.

Group	Criteria	Measurement	Measurement	Measurement	Measurement	Total
		A	B	C	D	
Mothers	Average	6.15	7.64	6.86	6.2	26.85
Normal children	Standard Deviation	3.38	3.28	2.58	3.64	10.40
Mothers	Average	5.05	4.58	4.5	3.59	17.71
Children With ADHD Disorder	Standard Deviation	3.14	2.89	2.88	2.85	8.40
Total	Average	5.6	6.11	5.68	4.89	22.28
	Standard Deviation	3.3	3.44	2.97	3.51	10.48

In this study the average age of most participants was between 36 to 42 years, the average age difference of mothers with normal children and mothers with hyperactive children was not noticeable. The average score of psychological health for most participants

stood from 12-33. In scale **A**, the average score of most mothers, almost was from 2-9. In scale **B**, the average scores were mostly 3-10. In scale **C**, most participants got a score from 2.3-8.6. Finally, in scale **D**, the average score was from 1.4-8.4. Average of mothers

with normal children in this five health measurements are obviously more than the average of mothers with ADHD Children.

Table 5- Results of **T-test** for comparison of two independent groups' mental health

Group	Lewin result's test in case of uniformity of variances			Uniformity of average T - test			
	P	F	Standard Error	Difference of Average	p	Degree of Freedom	t
Scale of A of mother's health psychology	0.32	1.01	0.52	1.10	0.04	158	2.13
Scale of B of mother's health psychology	0.35	0.89	0.49	3.06	0.00	158	6.27
Scale of C of mother's health psychology	0.69	0.16	0.43	2.36	0.000	158	5.46
Scale of D of mother's health psychology	0.05	3.86	0.52	2.61	0.000	158	5.06
Total	0.06	3.63	1.50	9.14	0.000	158	6.11

Table 5, shows the results of T-test for comparison of two independent groups' mental health; mothers with normal children and mother's with hyperactive children. Based on this table, the meaningful level of *Lewin* is set at $p > 0.05$. The results of table indicate that participants of the two groups were equal in their variance of psychological health and its subscales, but there is a significantly meaningful difference between mothers with normal children and ADHD children's mothers ($p < 0.05$).

With respect to the meaningful results and by paying attention to the results of Table

4, it can be said that level of mental health is significantly higher in mothers with normal children than that of mothers with ADHD children.

Discussion

Descriptive findings concerning age and psychological health of participants, as well as results of T-test were presented to compare the two independents groups (mothers with normal children and mothers with attention deficit / hyperactive disorder children).

The outcomes of the current study were in the same path with those kimiyae and beige (2010), Sanaei Kamal and college (2013), Beirami and college (2008), Shure and

Gaue(2007 Quoted By Sanaei Kamal, 2013) Hekmati and college(2009). These studies have shown that mothers and parents of children with attention deficit / hyperactivity disorder had lower levels of mental health than mothers of normal children.

The results of Chi-square test have also indicated that participants' scores varied across different educational levels. That is the frequency of participants was much higher at the first and second grades. This outcome is then similar to those achieved by Namdari and colleagues (2008).

Conclusion

As the literature and now the current study showed, parents' stress in families with attention deficit / hyperactivity's children is higher which consequently reduces the level of parents' psychological health. Further, considering Iranian cultural and social affairs, mothers spend more time with children than fathers. This maternal super visional role in children's education and social life causes many difficulties, especially for mothers with hyperactivity disorder / attention deficit children. Since these children need more devotion of time and effort to manage their duties and educational works, they are socially and educationally faced with various difficulties and challenges. Therefore, mothers dealing with this type of children are more prone to psychological damages, because they are constantly involved with their children. As

a result, this problem, if not dealt with properly, may end up in collapse of the structure and reduction of life-quality in families. Akhavan Karbasi et al. ^[39] reported that the spread rate of 16.3 for this disorder among 6-year old children in Yazd city is considered acceptable. Thus, it is essential to pay more attention to this disorder and try to reduce the number of these children and their families.

In a study conducted by Najafi et al., ^[40] it was presented that attention deficit/hyperactivity disorder is the first disorder among elementary-school children in Iran. Consequently, conductance of comprehensive programs is considered essential to interfere and educate parents (especially mothers) and teachers to reduce the individual and social damages. On the other hand, with attention to the results of researches based on abundance of this disorder in foundational education, it is proposed to assess this disorder at first, then, parents require to participate in special educational classes, and finally further interventions should be conducted at proper time to reduce educational and psychological damages in the future.

Acknowledgment

None .

Conflict of Interest

None .

References

- 1- Chuang LY, Tsai YJ, Chang YK, et al. Effects of acute aerobic exercise on response preparation in a Go/No Go Task in children with ADHD: An ERP study. *Journal of Sport and Health Science*. 2015; 4(1): 82- 88
- 2- Smith TF, Schmidt-Kastner R, McGeary JE, et al. Pre- and Perinatal Ischemia-Hypoxia, the Ischemia-Hypoxia Response Pathway, and ADHD Risk. *Behavior Genetic*. 2016; 46(3): 467-477
- 3- Esmaeilzadeh M, Qaderpour F, Ahmadi A. Interactive Vulnerability among Families having Children with ADHD: a Qualitative Study. *International Journal of Academic Research in Applied Science*.2014; 3(5):34- 43
- 4- Abedi A, Jamali S, Faramarzi S, et al. A comprehensive meta-analysis of the common interventions in ADHD. *Contemporary Psychology*. 2012; 7(1): 17-34
- 5- Bonvicini C, Faraone SV, Scassellati C. Attention-deficit hyperactivity disorder in adults: A systematic review and meta-analysis of genetic, pharmacogenetic and biochemical studies. *Molecular Psychiatry*. 2016; 21: 872–884
- 6- Piepmeier TA, Shih Ch H, Margaret W, et al. The effect of acute exercise on cognitive performance in children with and without ADHD. *Journal of Sport and Health Science*. 2015; 4(1):97- 104
- 7- Danforth JS, Harvey E, Ulaszek WR, et al. The outcome of group parent training for families of children with attention-deficit hyperactivity disorder and defiant/aggressive behavior. *Journal of Behavior Therapy and Experimental Psychiatry*. 2006; 37(3): 188-205
- 8- Mishra J, Sagar R, Joseph A, et al. Training sensory signal-to-noise resolution in children with ADHD in a global mental health setting. *Transl Psychiatry*.2016; 45:1- 9
- 9- García Murillo L, Ramos-Olazagasti M, Mannuzza S, et al. Childhood Attention-Deficit/Hyperactivity Disorder and Homelessness: A 33-Year Follow-Up Study. *Journal of the American Academy of Child & Adolescent Psychiatry* 2016; 55(11): 931–936
- 10- Sarraf N, Yousefi Kh, Majlesi F, et al. Determining effectiveness of multimodal parent management training combined with methylphenidate on academic achievement, oppositional and behavioral symptoms of school aged children with attention deficit hyperactivity disorder. *Razi Journal of Medical Sciences*.2015; 22(138): 32- 44. [Persian]
- 11- Alizadeh H, Hossein zadeh A, Godarzi AM, et al. Comparison of graphical patterns, ability of organizing and visual memory – spatial of children with or without attention deficit / hyperactivity. *Scientific Journal of Cognitive and Behavioral Sciences*. 2013; 3 (1): 1-10
- 12- Najarzadegan M, Nejati V, Amiri N, et al. Effect of cognitive rehabilitation on executive function (working memory and attention) in children with Attention Deficit Hyperactivity Disorder. *Journal of Rehabilitation Medicine*. 2014; 4(2): 97-108.
- 13- Hosseinkhanzadeh AA, Training social skills to children with attention deficit /hyperactivity disorder. *Exceptional education*. 2013; 13(8): 23- 41
- 14- Davari R. The Effectiveness of Integrated Cognitive-Behavior Dynamic Family - Oriented Treatment on Clinical and Executive Functions Symptoms in Children with Attention Deficit Hyperactivity Disorder. *Journal of Thought & Behavior in Clinical Psychology*. 2015; 9 (36): 67-76

- 15- Catala-Lopez F, Hutton B, Nunez-Beltran A, et al. The pharmacological and non-pharmacological treatment of attention deficit hyperactivity disorder in children and adolescents: protocol for a systematic review and network meta-analysis of randomized controlled trials. *Bio Med Central*. 2015; 4(19): 1- 10.
- 16- Gheragh Molaii L, Khosravi Z, BaniJamali S. The Role of Monitoring in Reducing Behavioral Problems in 6-12 Years Old Girls with ADHD: A Comparison between Maternal Monitoring and Self Monitoring. *Journal of Family Research*. 2014; 9(4):451-470
- 17- Weisani M, Shehni M, Alipour S, et al. Effectiveness of child-parent relation therapy on the symptoms of attention deficit / hyperactivity disorder (ADHD). *Family Psychology*.2015; 2(1): 29-38
- 18- Moosavi K, AmiriMajd M, Bazzazian S. A Comparison of Theory of Mind between Children with Autism, Intellectual Disability, ADHD, and Normal Children. *Middle Eastern Journal of Disability Studies*. 2014; 4(8):43-51.[Persian]
- 19- Kordlou M. hyperactivity: proper methods in relation with hyperactive children. *Journal of exceptional education*. 2015; 4(113): 38-48.
- 20- Kouchakzadeh S, Namazi A, Zarkesh M. Symptoms of attention deficit hyperactive disorder in children under school age according to the parent's information. *Nursing and Midwife magazine*. 2015 ;(25) 75: 35- 44. [Persian]
- 21- Kordestani, D. The effect of behavioral parent training on improving the mental health of mothers with attention deficit hyperactivity disorder children and decreasing their children's externalizing behavior. *Journal of Behavioral Sciences*. 2014; 8(3): 279-286
- 22- Jafari B, Fathi Ashtiyani A, Tahmasiyan K. Positive effects from group training program on mothers caused to reduce symptoms of attention deficit hyperactive disorder. *Journal of Counseling and Psychotherapy culture*. 2014; 7: 123-139.
- 23- Kimiaee SA, Baygi F. Comparison the family function of normal and ADHD mothers. *Journal of Behavioral Sciences*. 2010; 4(2): 141-147
- 24- Vaziri Sh, Lotfi Kashani F, Sorati M. Effectiveness of family training in reduced symptoms of the children with Attention Deficit Hyperactivity Disorder. *Procedia- Social and Behavioral Sciences* 2014; 128: 337– 342
- 25- Abdolkarini M, Mousavi VA, Khosrojauid M, et al. Effects on parents with behavioral management training on marriage satisfaction level of mothers with attention deficit hyperactive disorder. *Journal of Medical Sciences in Guilan University*. 2012; 20(80):74 -84
- 26- Behiyat S. Attention Deficit Hyperactivity Disorder Treatment Review; *CATALYST*. 2011; 6(1):12-17
- 27- Hakim javadi M, Lavasanai M, Shakoori, HR, et al. Comparison on effects of behavioral treatment, medicational treatment and the methods and the mixed methods of behavioral and medicational treatment on reducing symptoms of attention deficit hyperactivity disorder. *Medical science Journal of Gorgan*.2015; 53: 42-47.[Persian]
- 28- Mayes SD, Calhoun SL, Crowell EW. Learning disabilities and ADHD: overlapping spectrum disorders. *J Learn Disabil*.2000; 33: 417-424.

- 29- Gualtieri CT, Johnson LG. Efficient allocation of attentional resources in patients with ADHD: Maturational changes from age 10 to 29. *J AttenDisord.*2006; 9: 1-9.
- 30- Bahmani T, Alizadeh H. Study of effects on mothers behavioral management training in order to improve better child grows and reducing symptoms of attention deficit hyperactive disorder. *Special Individual Psychology.* 2011; 1: 27-39.
- 31- Sanaei Kamal S, Babapor J, Abdol Mohamadi K, et al. Comparing the Mental Health of Children's' Mothers With ADHD and Normal Children' Mothers in Hamedan. *European Online Journal of Natural and Social Sciences.* 2013;2(2):449-460
- 32- Beyrami M, Hekmati E, Soudmand M. Comparing performance families of children with ADHD disorder to families normal children, *Journal science-research Psychology of University of Tabriz* .2009; 4(15):21-41.[Persian]
- 33- Shure S, Gaue F. Parental and family factors for attention-deficit hyperactivity disorder in Taiwanese children. *Australian and New Zealand Journal of Psychiatry.* 2007; 41(8): 688-696
- 34- Hekmati E, Sodmand M, Beirami M, et al. Compare marital conflicts mothers of children with attention deficiencies disorder /hyperactivity with mothers of normal children. *Woman and Family Studies.* 2009;1(3):33-59.[Persian]
- 35- Abedi S, PourmohamadReza M, Mohamadkhani P, et al. Effectiveness of positive parenting program on mother-child relationship in children with attention deficit/ hyperactivity disorder. *Journal of Clinical Psychology.* 2012; 4(3): 63-73
- 36- Zargarinejad G. Efficacy of parent's training on problem behaviors in ADHD children. *Psychological Studies.*2007; 3(2): 29-48. [Persian]
- 37- Shafaat A, Tiregi A. Research of spread of anxiety symptoms in mothers with attention deficit hyperactive disorder, *Journal of Mazandaran, Medical Science.* 2010; 21(1) 66-72 [Persian]
- 38- Fathi Ashtiyani A. *psychological tests.* 4th ed, Tehran: Besat; 2010. [Persian]
- 39- Akhvan Karbasi S, Golestan M, Falah R, et al. The prevalence of attention deficit / hyperactivity disorder in children of 6 years in Yazd city. *Journal of Research in Medical Sciences of Shahid Sadoughi in Yazd.* 2008; 15(4): 29 –34 [Persian]
- 40- Najafi M, Fouladchang M, Alizadeh H, et al. Attention deficit disorder, conduct disorder and oppositional defiant disorder in primary school children. *Research on Exceptional Children.*2009; 3: 239-254