Failure to Thrive and its Risk Factors in 0-24 Months Children in Bojnurd City of Iran during 2008-2013

Hossein Lashkardoost ¹ , Saeid Doaei ² , Zohreh Akbari ³ , Fatemeh Mashkooti ³ , Ebrahim Hosseinzadeh ⁴ , Andishe Hamedi *⁵

- 1. Department of Epidemiology & Biostatistics, School of Public Health, North Khorasan University of Medical Sciences, Bojnurd, Iran
- 2. Department of nutrition sciences, Guilan University of Medical Sciences, Rasht, Iran
- 3. Mashhad University of Medical Sciences, Mashhad, Iran
- 4. Department of Pediatrics Nursing, North Khorasan University of Medical Sciences, Bojnurd, Iran
- 5. Faculty of nursing, Shirvan Center of Higher Health Education, North Khorasan University of Medical Sciences, Bojnurd, Iran

ARTICLE INFO

Original Article

Received: 21Sep 2019 Accepted: 20Jan 2020



Corresponding Author:

Andishe Hamedi ahamedi1364@gmail.com

ABSTRACT

Introduction: Failure to thrive (FTT) is a global problem and one of the most common health problems in childhood that involves many other social, economic, and cultural factors. Considering the adverse effects of FTT in the future of children, we studied FTT and its related factor in children under the age of 2 years in Bojnurd (the capital city of North Khorasan province, Iran).

Methods: This study was a Retrospective cohort study on 1000 health records, born in 2008-2013. Stratified sampling method was applied and the data were collected using a checklist in the health centers. Finally, data were analyzed using Chi-square, Multiple logistic regression, and independent t-test in SPSS₁₉ software. Significant level was set at 5%.

Results: Incidence of FTT was calculated as 443 children (44.3%) in the children's first two years of life. A significant relationship was observed between FTT in children and head circumference disorders at birth (p=0.001), maternal age at delivery (p=0.01), mother's education level (OR=0.4 CI95% [0.2-0.8] p=0.012), type of delivery (OR=0.5 CI95% [0.4-0.7] p<0.001), unspecified gestational age (OR=3.6 CI95% [1.3-10.08 p=0.015]), and pregnancy under the age of 18 (OR=2.4 CI95% [1.1-5.3] p=0.02).

Conclusion: Considering the high incidence of FTT in children, increasing awareness about timely feeding, promoting households' health, preventing and controlling infectious diseases should be improved.

Keywords: Failure to Thrive, Risk Factors, Children

How to cite this paper:

Lashkardoost H, Doaei S, Akbari Z, Mashkooti F, Hosseinzadeh E, Hamedi A. Failure to Thrive and its Risk Factors in 0-24 Months Children in Bojnurd City of Iran during 2008-2013. J Community Health Research. 2020; 9(1): 13-20.

Copyright: ©2020 The Author(s); Published by Shahid Sadoughi University of Medical Sciences. This is an open-access article distributed under the terms of the Creative Commons Attribution License (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Failure to thrive (FTT) is a global problem and one of the most common health problems in childhood that involves many other social, economic, and cultural factors (1). In fact, FTT is associated with an inadequate physical growth or inability to maintain the desired growth rate over time in children (2). In developing countries, FTT causes disrupt in children's weight gain, mental development, mental health, and defense against infection. According to WHO, more than 30% of children under the age of 5 years suffer from FTT, of which 80% have stunting and 20% are underweight (3, 4).

According to the related studies, the prevalence of FTT in third world countries is higher than other parts of the world. In most of these countries, the physical growth of children is lower than the international standard. In our country 16.3% of girls, 15% of boys, and totally 15.7% of children are underweight and 18.4% of girls, 19.5% of boys, and totally 18.9% of children are stunting (5). In Iran, the most prevalent ages of FTT and stunting are in 6-12 and 12-24 months, respectively. The former is the time of initiating supplementary feeding (6). Low birth weight increases with age, so that at 2 years of age it reaches a peak of 13.8% (3).

One of the most important causes of FTT is deprivation of the baby from breast feeding, early or late start of the supplementary feeding, not giving enough food to the child, child illness, emotional and psychological problems, non-compliance with health issues, and use of unhealthy water(7).

The studies showed that 46% of the FTT were due to non-specific causes (poverty and lack of access to inadequate food, low parenting education) and 26% of the cases were due to organ causes (5). Obviously, identifying risk factors is possible by monitoring the growth of children. Growth monitoring is one of the most important sources of information to diagnose FTT and malnutrition in children.

Moreover, FTT is usually a very slow process, not recognizable by families. Along with the above-mentioned issues, adverse effects of future growth disorder in children, include possible mortality, increased related diseases, reduced learning ability, and reduced mental and physical capacity. Considering few studies on the risk factors of FTT conducted in Iran, we decided to examine FTT in children under 2 years of age and its effective factors in the city of Bojnurd.

Methods

Study design: The present research was a retrospective cohort study on 1000 health records. The target population was children who were born in Bojnurd City and had health records in the years of 2008-2013.

Sampling and sample size

Stratified sampling method (based on the year of birth and urban sanitary facility in Bojnurd) was conducted and the quota of each center was determined. Based on simple random sampling, 1000 health records were selected. Sample size was determined as 1000 people according to the results reported by Naderi (8) considering the expected prevalence of malnutrition at about 5%, 95% confidence level, and accuracy of 0.01.

The inclusion criteria in this research were completeness of the child' health file in terms of the variables under consideration. In the case of incomplete data (due to being illegible, incomplete, confusing, etc.), the following file was used for review.

Instrument and data gathering

The data needed in this study were collected using a checklist derived from maternal and child care records available in health centers. Information collected from this checklist included the physical growth of children (weight, height, head circumference) over two years. These data of children were extracted from their household health records. The FTT was determined in children whose information with regard to these items was not parallel to or had an ascending trend considering the previous growth Furthermore, information such as demographic characteristics and underlying diseases of the mother as well as the previous pregnancy status of mothers were collected. Finally, data were analyzed by Chi-square, Multiple logistic regression, and independent t-test using SPSS₁₉ software. Significant level was considered as 0.05.

Results

In this study, 1000 health records of children were studied; 522 of them were boys (52.2%). The mean and standard deviation of the maternal age at delivery was 26.8 ± 5.3 years. We found that 87.5% of mothers were housewives and 44.3% had undergraduate education. Based on the results,

incidence of FTT occurred in 443 children (44.3%) in the first two years of life, while 557 children had no FTT (55.7%). About 146 (15%) children were stunting and 368 (37.5%) had weight FTT. The relative frequencies of FTT are shown in Fig 1.

The highest FTT in 18 months of age was 13.2%, the least FTT 3-5 after birth was 0.1%, and no abnormality was observed at birth. The relative frequencies of FTT based on the months are represented in Fig. 2.




Figure 1. Relative frequency of Failure to thrive.




Figure 2. Relative frequency of Failure to thrive based on Months of the Year

The quantitative characteristics of the factors affecting FTT are presented in Table 1.

As show in Table 1, a significant difference was

found between maternal age at delivery and head circumference in children with FTT in two groups (p<0.05).

Table 1. Characteristics of quantitative variables which affecting Failure to thrive

Variable	of children without FTT (Mean±SD)	of children with FTT (Mean±SD)	P-value
Birth weight	$3.4{\pm}1.6$	3.2±0.4	0.06
Height at birth	50.8 ± 4.1	50.5±3.6	0.35
Head circumference at birth	34.5±1.6	30.01 ± 1.8	0.001^{*}
Maternal age at delivery	27.2±5.3	26.3±5.3	0.01^{*}
Mother BMI	24.8 ± 4.3	25.02 ± 4.6	0.52
Child birth rank	1.8 ± 0.9	1.8 ± 0.8	0.58
Time to start auxiliary feeding	6.2 ± 0.6	6.2 ± 0.6	0.69

^{*}Significant at 0.05% level

The FTT was higher in children whose mother's age was under 18 years. Furthermore, FTT was lower in children whose mothers had higher education levels. A significant relationship was observed between type of delivery and unspecified

gestational age with FTT in children (p<0.05). However, no significant relationship was found between mother's history of previous pregnancy and FTT in children (p=0.45) (Table 2).

Table 2. Odds ratios and confidence intervals of qualitative variables which affecting Failure to thrive

Variabl	es	children without FTT	children with FTT	Crude OR and 95% CI **	P-value	Adjusted OR and 95%CI ¶	P- value
Gender	Male Female	289(51.9) 268(48.1)	233(52.5) 210(47.5)	1 0.9(0.7-1.2)	0.84		
Mother's job	housewife Employee	480(86.6) 77(13.4)	398(90.2) 45(9.8)	1 0.6(0.4-1.03)	0.07	1 0.7(0.44-1.2)	0.27
Mother's education level	Illiterate Under diploma Academic	28(4.8) 177(31.9) 352(63.3)	31(7.1) 207(46.6) 205(46.3)	1 0.9(0.5-1.7) 0.4(0.2-0.8)	0.96 0.012*	1 0.6(0.35-1.1) 0.6(0.3-1.3)	0.23 0.04
Mother's height less than 150cm	No Yes	538(96.6) 19(3.4)	430(97.3) 13(2.7)	1 0.7(0.3-1.6)	0.52	0.0(0.3-1.3)	0.04
Type of delivery	Cesarean Natural	294(52.7) 263(47.3)	291(65.8) 152(34.2)	1 0.5(0.4-0.7)	<0.001*	1 0.7(0.54-1.02)	0.001
Number of children	<=2 2-5 >=5	414(74.4) 133(23.9) 10(1.6)	319(72.3) 114(25.7) 10(2)	1 1.1(0.8-1.4) 1.2(0.5-3.2)	0.5 0.6		
First pregnancy	No Yes	317(56.9) 240(43.1)	265(60) 178(40)	1 0.8(0.6-1.1)	0.31		
Fifth and above pregnancy	No Yes	537(96.4) 20(3.6)	426(96.2) 17(3.8)	1 1.07(0.5-2.07)	0.83		
Pregnancy under the age of 18	No Yes	410(98.2) 147(1.8)	424(95.7) 19(4.3)	1 2.4(1.1-5.3)	0.02*	1 2.1(0.9-4.6)	0.01
Pregnancy over 35	No	523(93.9)	420(95)	1			

^{**}Independent t-test

years old	Yes	34(6.1)	23(5)	0.8(0.4-1.3)	0.43		
Pregnancy interval less than 3 years	No Yes	520(93.4) 37(6.6)	410(92.8) 33(7.2)	1 1.09(0.6-1.7)	0.71		
Multidimensional	No Yes	542(97.6) 15(2.4)	433(97.5) 10(2.5)	1 1.05(0.4-2.3)	0.89		
Baby weight less than 2500 kg	No Yes	546(98) 11(2)	441(99.5) 2(0.5)	1 0.2(0.05-1.02)	0.05	1 0.2(0.04-1.01)	0.23
Baby weight over than 4000 kg	No Yes	552(99.1) 5(0.9)	439(99.1) 4(0.9)	1 1.006(0.2-3.7)	0.99		
	Breast	483(86.7)	395(89.2)	1			
Type of infant feeding	feeding Synthetic milk	14(2.5)	9(2.1)	0.7(0.3-1.8)	0.58		
	Both	60(10.7)	39(8.7)	0.7(0.5-1.2)	0.28		
Mother's desirable weight gain during	No Yes	141(26.6) 416(73.4)	122(29.4) 321(70.6)	1 0.8(0.6-1.1)	0.23		
mother's disease history	No Yes No	483(86.7) 74(13.3) 543(97.7)	365(82.6) 78(17.4) 433(97.7)	1 1.3(0.9-1.9) 1	0.07	1 1.1(0.6-2.1)	0.33
Risky behavior in mother or spouse	Yes	14(12.3)	10(2.3)	0.9(0.4-2.2)	0.93		
The history of high risk pregnancy	No Yes	420(75.4) 137(24.6)	343(77.4) 100(22.6)	1 0.8(0.6-1.2)	0.45		
Unspecified gestational age	No Yes	552(99.1) 5(0.9)	429(96.8) 14(3.2)	1 3.6(1.3-10.08)	0.015*	1 1.4(0.8-2.2)	0.04
unwanted pregnancy	No Yes	496(89) 61(11)	400(90.3) 43(9.7)	1 0.8(0.5-1.3)	0.52		
RH negative	No Yes	526(94.4) 31(5.6)	414(93.5) 29(6.5)	1 1.1(0.7-2.004)	0.51		
History of abortion	No Yes	542(97.3) 15(2.7)	426(95.9) 17(4.1)	1 1.5(0.7-3.07)	0.23		
Still birth	No Yes	549(98.6) 8(1.4)	436(98.4) 7(1.6)	1 1.1(0.3-3.06)	0.85		
Preeclampsia	No Yes	544(97.7) 13(2.3)	431(97.3) 12(2.7)	1 1.1(0.5-2.5)	0.70		
Infertility history	No Yes	546(98) 11(2)	433(97.7) 10(2.3)	1 1.1(0.4-2.7)	0.75		

^{*} Significant at 0.05% level

¶ Multiple logistic regression

Discussion

In the present study, the incidence of FTT in children's first two years of life was 44.3 with a confidence interval of 41.1-47.4. Of these children, 37.4% were underweight and 15% were stunting. Similar studies reported that low birth weight among children was about 26.6%(6). In this study, children's FTT started from their first year of life

and its highest rate was about 13.2% at 18 months. In addition, a significant relationship was found between head circumference at birth and FTT in children; children with FTT had less mean head circumference. Study conducted by Pronman et al. confirmed these results. They confirmed that children aged 0-11 months were less likely to develop FTT, which may be due to the protective

^{**} Chi-square, Univariate logistic regression

role of the breast feeding (4). At 12 months of age, a further FTT appears, which can be attributed to inappropriate complementary nutrition (9).

In the present study, no significant relationship was found between the history of high risk pregnancy in mothers and FTT. Study conducted by Blair and Hvelplund reported different results. So, history of abortion, stillbirth, and cesarean section had a significant relationship with children's FTT(10, 11). A significant relationship was also reported between maternal age during labor and FTT in children. Mothers who were younger than 18 years at the time of delivery were more likely to develop FTT in their children. However, no significant relationship was found in mothers over than 35 years of age. Hien showed that the risk of developing child's FTT increased in mothers under 24 years and over 35 years. This can be attributed to the mothers' lack of readiness for taking care of their children or their boredom (6).

No significant relationship was found between birth ranking, number of children, first pregnancy, fifth pregnancy, and FTT, but research conducted by Hein, Hvelplund, and Mohammadpoorasl showed that children born in families with more than 3 to 5 children were more likely to have FTT. This can be justified by the fact that children were less considered in such families by parents (6, 11, 12). In this study, we observed no significant relationship between the gestational interval less than 3 years and FTT in children, but Nahar and Victora indicated that less than 3 years of interval between pregnancies increased the risk of developing FTT in children, which contradicts with our study (13, 14).

In this study, maternal BMI, height less than 150 cm, and mother's desirable weight gain during pregnancy had no significant relationship with FTT in children. Emond et al.'s results contradicted with our in this regard. They mentioned that maternal weight and height played an important role in the development of FTT (15).

No significant relationship was also found between the type of infant feeding and the onset of supplemental nutrition with FTT in children. However, such studies indicated that long-term breastfeeding, early or later onset of supplemental nutrition, increased the incidence of FTT in children (1, 16). The current nutrition recommendations provided by Pediatric the Academy include exclusive breastfeeding for almost the first six months after birth. The best time to start complementary feeding is after 6 months of age, because at the end of 6 months of age, breast milk alone could not supply the needs of the child. Additionally, the gastrointestinal tract after 6 months of age has an admission for semisolid foods. If the supplemental nutrition starts earlier than this time, the risk of diarrhea and FTT would increase due to the lack of evolution of gastrointestinal tract.

In addition, FTT had no significant relationship with the child's gender and mother's occupation. However, Radhakrishna et al. reported that boys (17) were more susceptible to FTT than girls (11) and the incidence of FTT was greater in working mothers. These findings contradict with the results of this study (7, 18).

In the present study, a significant relationship was found between mother's education level and FTT in children, so that having higher education had a preventive role in the development of children's FTT. Nahar and Khuwaja confirmed these results, which can be attributed to participants' better management, greater use of health services, and promotion of health care (13, 19). However, such studies did not show a significant relationship between the level of mothers' education and childhood FTT(3). In the present study, a significant relationship was found between the type of delivery and FTT. Natural delivery was a preventative factor in the development of childhood FTT; this result was confirmed by a study (11). In this study, a significant relationship was found between uncertain pregnancy age and the incidence of childhood FTT; this finding was confirmed by a research (20). We found no significant relationship between mother's disease history and FTT, which is contrary to a study (10).

Mother's high risk behavioral factors had no significant relation with childhood FTT, which can

be attributed to low maternal reporting due to their social shame. However, a study on risk factors for weight faltering in infancy according to age at onset reported significant correlation between mother's high risk behavioral factors and childhood FTT (21).

Limitation

Some data were corrupted due to the filing and could not be accessed. Furthermore, the data were not recorded for research purposes, which restricted our findings.

As we did not evaluate dietary intakes and dietary supplements, future researchers are recommended to evaluate the relationship of dietary intakes and dietary supplements with growth retardation.

Conclusion

The results of this study showed no significant

relationship between high risk behavioral factors in mothers and childhood FTT. Regarding the high incidence of FTT in children, increasing parents' awareness, starting the nutritional supplements timely, promoting household health, as well as preventing and controlling infectious diseases are recommended to improve the growth of children.

Conflict of Interest

The authors declared that they have no conflict of interest.

Acknowledgements

Ethics Committee of North Khorasan University of Medical Sciences was: IR.NKUMS.REC.1394.839

The authors would like to appreciate the health center staff who helped us in conducting the present study.

Reference

- 1. Wright CM, Parkinson KN, Drewett RF. The influence of maternal socioeconomic and emotional factors on infant weight gain and weight faltering (failure to thrive): data from a prospective birth cohort. Archives of disease in childhood. 2006; 91(4):312-7.
- 2. Berak M, Azari Namin L, Nemati A, et al. Failure to Thrive Risk Factors among Infants in Namin. Journal of Ardabil University of Medical Sciences. 2003; 3(4):14-18.[Persian]
- 3. Gohari MR, Salehi M, Zaeri F, et al. Application of random effect model for determining factors affecting FTT in less than 2 years children in east of Tehran. Razi Journal of Medical Sciences. 2012; 19(99):32-9.[Persian]
- 4. Pronman L, Rondinelli M, Burkardt DD, et al. Pearson Syndrome: A Rare Cause of Failure to Thrive in Infants. Clinical pediatrics. 2019; 58(7):819-24.
- 5. Shekhi MF, Shamsi M, Khorsandi M, et al. Predictors accident structures in mothers with children under 5 years old in city of Khorramabad based on Health Belief Model. Safety Promotion and Injury Prevention. 2015; 3(3):199-206. [Persian]
- 6. Hien NN, Kam S. Nutritional status and the characteristics related to malnutrition in children under five years of age in Nghean, Vietnam. Journal of Prevention Medical Public Health. 2008; 41(4):232-40.
- 7. AM Shamsir A, Ahmed T, Roy S, et al. Determinants of Under Nutrition in Children under 2 years of age from Rural Bangladesh. Indian Academy of Pediatrics. 2012; 49.
- 8. Naderi S. Growth indices of 2-5 years old infants in Bandar Abbas day-care centers. Hormozgan Medical Journal. 2003; 6(4):31-35.[Persian]
- 9. Nojoumi M, Tehrani A, Najmabadi S. Risk analysis of growth failure in under-5-year children. 2004; 7(3):195-200.
- 10. Blair PS, Drewett RF, Emmett PM, et al. Family, socioeconomic and prenatal factors associated with failure to thrive in the Avon Longitudinal Study of Parents and Children (ALSPAC). International journal of epidemiology. 2004; 33(4):839-47.
- 11. Hvelplund C, Hansen BM, Koch SV, et al. Perinatal Risk Factors for Feeding and Eating Disorders in Children Aged 0 to 3 Years.Official Journal of the American Academy of Pediatrics. 2016; 137(2):2015-2575.
- 12. Mohammadpoorasl A, Sahebihag M, Rostami F, et al. Factors related to undesirable growth of 6 month–2years old children in Tabriz-Iran. Journal of gorgan university of medical sciences. 2010; 12(3):45-50.
- 13. Nahar B, Ahmed T, Brown KH, et al. Risk factors associated with severe underweight among young children reporting to a diarrhoea treatment facility in Bangladesh. Journal of health, population, and nutrition. 2010; 28(5):476.

- 14. Victora CG, Adair L, Fall C, et al. Maternal and child undernutrition: consequences for adult health and human capital. The lancet. 2008; 371(9609):340-57.
- 15. Emond A, Drewett R, Blair P, et al. Postnatal factors associated with failure to thrive in term infants in the Avon Longitudinal Study of Parents and Children. Archives of disease in childhood. 2007; 92(2):115-9.
- 16. Daymont C, Hoffman N, Schaefer E, et al. Clinician diagnoses of failure to thrive before and after switch to World Health Organization growth curves. Academic pediatrics. 2019.
- 17. Radhakrishna R, Ravi C. Malnutrition in India: Trends and determinants. Economic and Political Weekly. 2004; 39(7):671-6.
- 18. Khalil HM, Husain SM, Alkhateeb MS, et al. Failure To Thrive In Children Under Two Years Of Age And Associated Factors, A Hospital-Based Study. Diyala Journal of Medicine. 2019; 17(2):127-35.
- 19. Khuwaja S, Selwyn BJ, Shah SM. Prevalence and correlates of stunting among primary school children in rural areas of southern Pakistan. Journal of tropical pediatrics. 2005; 51(2):72-7.
- 20. Goh LH, How CH, Ng KH. Failure to thrive in babies and toddlers. Singapore medical journal. 2016; 57(6):287.
- 21. Olsen EM, Skovgaard AM, Weile B, et al. Risk factors for weight faltering in infancy according to age at onset. Paediatric and perinatal epidemiology. 2010; 24(4):370-82.